Corrigendum: Comparison of changes in the buccal keratinized mucosa between the minimally-invasive crestal incision and apically-positioned full-thickness flap methods for stage-two implant surgery

Keon-Il Yang¹, Jun-Mo Yang², Ki-Won Lee², Won-Pyo Lee³, Byung-Ock Kim⁴, and Sang-Joun Yu^{4*}

¹Clinical Professor, Department of Periodontology, Chosun University School of Dentistry, Gwangju, Republic of Korea

²Graduate Student, Department of Periodontology, Chosun University School of Dentistry, Gwangju, Republic of Korea

³Assistant Professor, Department of Periodontology, Chosun University School of Dentistry, Gwangju, Republic of Korea

⁴Professor, Department of Periodontology, Chosun University School of Dentistry, Gwangju, Republic of Korea

This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted noncommercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Correction to: Oral Biol Res 2020;44:45–50, <https://doi.org/10.21851/obr.44.01.202003.45>

In the originally published version of this article was error in the author's affiliation, which should be properly revised as follows:

Before correction:

Ki-Won Lee³

³Doctor of Dental Clinic, Samho Myeongin Clinic, Yeongam, Republic of Korea

After correction:

Ki-Won Lee²

²Graduate Student, Department of Periodontology, Chosun University School of Dentistry, Gwangju, Republic of Korea

***Corresponding author:** Sang-Joun Yu, Department of Periodontology, Chosun University School of Dentistry, 309 Pilmun-daero, Dong-gu, Gwangju 61452, Republic of Korea.

Tel: +82-62-220-3850, **Fax:** +82-62-224-4664, **E-mail:** sjyu78@chosun.ac.kr

Copyright © 2021, Oral Biology Research Institute